

Supplementary Information

Etanercept-Synthesising Mesenchymal Stem Cells Efficiently Ameliorate Collagen-Induced Arthritis.

Authors: Narae Park¹, Yeri Alice Rim¹, Hyerin Jung¹, Juryun Kim¹, Hyoju Yi^{1,2}, Youngkyun Kim^{1,2}, Yeonsue Jang¹, Seung Min Jung¹, Jennifer Lee¹, Seung-Ki Kwok¹, Sung-Hwan Park¹, and Ji Hyeon Ju¹ *

Affiliations:

1. Division of Rheumatology, Department of Internal Medicine, Seoul St. Mary's Hospital, College of Medicine, The Catholic University of Korea, Seoul, 137-701, Republic of Korea.

2. Department of Medicine, Institute for Stem Cell Biology and Regenerative Medicine, and Stanford Cardiovascular Institute, Stanford University School of Medicine, Stanford, CA, USA.


***Corresponding author.**

Ji Hyeon Ju, M.D., Ph.D.


Division of Rheumatology, Department of Internal Medicine, Seoul St. Mary's Hospital, College of Medicine, The Catholic University of Korea, Banpodaero 222, Seochogu, Seoul, 137-701, Republic of Korea, Tel: 82-2-2258-6893, Fax: 82-2-3476-2274, E-mail: juji@catholic.ac.kr.

Supplementary figures

a



b



Supplementary Figure S1. Generation and drug expression of the mcTNFR2. (a) The full length gel electrophoresis of parental plasmids (mock and sTNFR2-Fc) after arabinose incubation. Minicircles with an appropriate size were detected after arabinose induction (arrows). (b) The full length western blot of sTNFR2-Fc expression. sTNFR2-Fc was detected in the conditioned media of mcTNFR2-transfected HEK293T cells (arrows). Etanercept was used as a control.